

Galois representations attached to étale cohomology

Sebastian Petersen

Universität der Bundeswehr, Neubiberg, Germany
sebastian.petersen@unibw.de

Session: 3. Arithmetic Geometry

Let K be a field and X/K a separated algebraic scheme. Grothendieck and Artin constructed for every prime number ℓ and every $i \in \mathbf{N}$ a cohomology group $H^i(X_{\overline{K}}, \mathbf{Q}_\ell)$ which comes with a natural action of the Galois group $\text{Gal}(\overline{K}/K)$; one thus obtains a Galois representation

$$\rho_\ell: \text{Gal}(\overline{K}/K) \rightarrow \text{Aut}_{\mathbf{Q}_\ell}(H^i(X_{\overline{K}}, \mathbf{Q}_\ell)).$$

Denote by $\rho: \text{Gal}(\overline{K}/K) \rightarrow \prod_\ell \text{im}(\rho_\ell)$ the homomorphism induced by the ρ_ℓ . Serre proved recently that in the case where K is a number field the family $(\rho_\ell)_\ell$ is almost independent in the following sense: There exists a finite Galois extension K'/K such that

$$\rho(\text{Gal}(\overline{K}/K')) = \prod_\ell \rho_\ell(\text{Gal}(\overline{K}/K')).$$

This information is quite useful when working with such families of ℓ -adic representations attached to schemes, and it ties in well with the adelic openness conjecture. There are analogous results in the case where K is an arbitrary finitely generated field of characteristic zero, and where K is a geometric function field of arbitrary characteristic. Following a suggestion of Illusie and making strong use of results of Orgogozo this was used to establish a quite general independence theorem for families $(F_\ell)_\ell$ of étale sheaves of \mathbf{F}_ℓ -vector spaces over an arithmetic scheme which satisfy a uniform constructability and a potential semistability condition.